

# One Health

Psico-neuro-endocrino-immunologia, psichiatria di liaison  
e problemi medici per la persona con disturbi del neurosviluppo



Mercoledì 3 Dicembre 2025





Facoltà Teologica dell'Italia Centrale, Firenze

Il contributo delle scienze umane alla liaison psichiatrica

**Roberto Franchini**, Segretario SIDiN, Università Cattolica del Sacro Cuore, Responsabile Area Strategica e Sviluppo Opera Don Orione Italia,

**Corrado Cappa**, Consigliere SIDiN, Psichiatra Dipartimento Salute Mentale e Dipendenze Patologiche AUSL Piacenza

*Il corpo escluso*





**DANZA IN CARROZZINA**

**e DanceAbility**

 **DISABILI**NEWS.COM






individualità

universalità

















# **COMING OUT PROUD**

per eliminare lo stigma della malattia mentale

## *WORKBOOK*


PER I FACILITATORI E I PARTECIPANTI AL PROGRAMMA

Blythe A. Buchholz and Patrick W. Cormican



Traduzione a cura di  
Giulia Reali  
Paola Carozza

Revisione dell'editing  
Lorenzo Benini



## Prescribing psychotropic drugs to adults with an intellectual disability

Julian N Trollor

**Table 1 Common medical comorbidities in people with intellectual disability that may alter the choice of psychotropic drug**

Comorbidity	Associated genetic syndromes	Prescription Implications
Epilepsy	Down, Fragile X, Angelman, Tuberous sclerosis, Rett, Wolf-Hirschhorn	Exercise caution prescribing psychotropics that lower seizure threshold, e.g. clozapine, tricyclic antidepressants, venlafaxine
Obesity	Down, Turner, Angelman, Prader-Willi	Avoid psychotropics with high cardiometabolic liability as first-line treatment
Dyslipidaemia	Down, Turner, Prader-Willi	Avoid psychotropics with high cardiometabolic liability as first-line treatment
Type 2 diabetes	Down, Turner, Sotos, Prader-Willi	Avoid psychotropics with high cardiometabolic liability as first-line treatment
Hypertension	Turner, Tuberous sclerosis, Williams, Sotos, Prader-Willi	Exercise caution prescribing psychotropics known to raise blood pressure, e.g. venlafaxine, desvenlafaxine, duloxetine
Hypotension	Down	Where possible avoid psychotropics with potential to exacerbate, e.g. chlorpromazine, tricyclic antidepressants, quetiapine
Respiratory difficulties or structural airway abnormalities	Prader-Willi, Down	Where possible avoid highly sedating psychotropics that may exacerbate the risk of respiratory failure
Swallowing difficulties	Cerebral palsy	Exercise caution with psychotropics that exacerbate swallowing difficulties, e.g. clozapine, olanzapine, risperidone, quetiapine
Early onset dementia	Down	Be aware that cognitive adverse effects of some psychotropics may compound cognitive dysfunction in emerging dementia









*Silenzio:  
nel dirupo inaccessible,  
al fondo,  
il prato in fiore*